



Perfect solution to all problems

Tips, Tricks, General Knowledge, Current Affairs, Latest Sample,
Previous Year, Practice Papers with solutions.

CBSE 12th Chemistry 2017 Unsolved Paper Delhi Board

Buy solution: <http://www.4ono.com/cbse-12th-chemistry-solved-previous-year-papers/>

Note

This pdf file is downloaded from www.4ono.com. Editing the content or publicizing this on any blog or website without the written permission of [Rewire Media](#) is punishable, the suffering will be decided under DMCA

CBSE 12th Chemistry 2017 Unsolved Paper

Delhi Board


TIME - 3HR. | QUESTIONS - 26

THE MARKS ARE MENTIONED ON EACH QUESTION

SECTION – A

Q.1. Write the formula of an oxo-anion of Manganese (Mn) in which it shows the oxidation state equal to its group number. 1 mark

Q.2. Write IUPAC name of the following compound: 1 mark





Q.3. For a reaction $\text{R} \rightarrow \text{P}$, half-life ($t_{1/2}$) is observed to be independent of the initial concentration of reactants. What is the order of reaction? 1 mark

Q.4. Write the structure of 1-Bromo-4-chlorobut-2-ene 1 mark

Q.5. Write one similarity between Physisorption and Chemisorption. 1 mark

SECTION - B

Q.6. Complete the following reactions: 2 marks



OR

What happens when

(i) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ is heated?

(ii) H_3PO_3 is heated?

Write the equations.

Q.7. Define the following terms: 2 marks

- (i) Colligative properties
- (ii) Molality (m)

Q.8. Draw the structures of the following: 2 marks

- (i) $\text{H}_2\text{S}_2\text{O}_7$
- (ii) XeF_6

Q.9. Calculate the degree of dissociation (α) of acetic acid if its molar conductivity (Λ_m) is

$39.05 \text{ S cm}^2 \text{ mol}^{-1}$. 2 marks

Given $\lambda^\circ(\text{H}^+) = 349.6 \text{ S cm}^2 \text{ mol}^{-1}$ and $\lambda^\circ(\text{CH}_3\text{COO}^-) = 40.9 \text{ S cm}^2 \text{ mol}^{-1}$

Q.10. Write the equations involved in the following reactions: 2 marks

- (i) Wolff-Kishner reduction
- (ii) Etard reaction

SECTION - C

Q.11. A 10% solution (by mass) of sucrose in water has freezing point of 269.15 K.

Calculate the freezing point of 10% glucose in water, if freezing point of pure water is 273.15 K. 3 marks

Given: (Molar mass of sucrose = 342 g mol^{-1})

(Molar mass of glucose = 180 g mol^{-1})

Q.12. (a) Calculate the mass of Ag deposited at cathode when a current of 2 amperes was passed through a solution of AgNO_3 for 15 minutes. 3 marks

(Given: Molar mass of Ag = 108 g mol^{-1} $1\text{F} = 96500 \text{ C mol}^{-1}$)

(b) Define fuel cell.

Q.13. (i) What type of isomerism is shown by the complex $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$?

(ii) Why a solution of $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ is green while a solution of $[\text{Ni}(\text{CN})_4]^{2-}$ is colourless? (At. no. of Ni = 28). 3 marks

(iii) Write the IUPAC name of the following complex: $[\text{Co}(\text{NH}_3)_5(\text{CO}_3)]\text{Cl}$.

Q.14. Write one difference in each of the following: 3 marks

- (i) Lyophobic sol and Lyophilic sol
- (ii) Solution and Colloid
- (iii) Homogeneous catalysis and Heterogeneous catalysis

Q.15. Following data are obtained for reaction: 3 marks



t/s	0	300	600
(N ₂ O ₅)/mol L ⁻¹	1.6×10^{-2}	0.8×10^{-2}	0.4×10^{-2}

(a) Show that it follows first order reaction.

(b) Calculate the half-life.

(Given $\log 2 = 0.3010$, $\log 4 = 0.6021$)

Q.16. Following compounds are given to you: 3 marks

2-Bromopentane, 2-Bromo-2-methylbutane, 1-Bromopentane



- (i) Write the compound which is most reactive towards S_N2 reaction.
- (ii) Write the compound which is optically active.
- (iii) Write the compound which is most reactive towards β-elimination reaction.

Q.17. (a) Write the principle of method used for the refining of germanium.

(b) Out of PbS and PbCO₃ (ores of lead), which one is concentrated by froth floatation process preferably? 3 marks

(c) What is the significance of leaching in the extraction of aluminium?

Q.18. Write structures of compounds A, B and C in each of the following reactions: 3 marks



OR

Do the following conversions in not more than two steps:

- (i) Benzoic acid to benzaldehyde
- (ii) Ethyl benzene to Benzoic acid
- (iii) Propanone to Propene

Q.19. Write the structure of the monomers used for getting the following polymers: 3 marks

- (i) Dacron
- (ii) Melamine – formaldehyde polymer
- (iii) Buna-N

Q.20. Define the following:

- (i) Anionic detergents
- (ii) Broad spectrum antibiotics
- (iii) Antiseptic

Q.21. Give reasons: 3 marks

- (i) Thermal stability decreases from H_2O to H_2Te .
- (ii) Fluoride ion has higher hydration enthalpy than chloride ion.
- (iii) Nitrogen does not form pentahalide.

Q.22. Give reasons: 3 marks

- (i) Acetylation of aniline reduces its activation effect.
- (ii) CH_3NH_2 is more basic than $\text{C}_6\text{H}_5\text{NH}_2$.
- (iii) Although $-\text{NH}_2$ is o/p directing group, yet aniline on nitration gives a significant amount of m-nitroaniline.

SECTION - D

Q.23. After watching a programme on TV about the presence of carcinogens (cancer causing agents) Potassium bromate and Potassium iodate in bread and other bakery products, Ritu a class XII student decided to aware others about the adverse effects of these carcinogens in foods. She consultanted the school principal and requested him to instruct canteen contractor to stop selling sandwiches, pizza, burgers and other bakery products to the students. Principal took an immediate action and instructed the canteen contractor to replace the bakery products with some proteins and vitamins rich food like fruits, salads, sprouts etc. The decision was welcomed by the parents and students. 4 marks

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Ritu?
- (ii) Which polysaccharide component of carbohydrates is commonly present in bread?
- (iii) Write the two types of secondary structure of proteins.
- (iv) Give two examples of water soluble vitamins.

SECTION – E

Q.24. (a) Account for the following: 5 marks

- (i) Transition metals form large number of complex compounds.
- (ii) The lowest oxide of transition metal is basic whereas the highest oxide is amphoteric or acidic.
- (iii) E° value for the Mn^{3+}/Mn^{2+} couple is highly positive (+1.57 V) as compare to Cr^{3+}/Cr^{2+} .
- (b) Write one similarity and one difference between the chemistry of lanthanoid and actinoid elements.

OR

- (a) (i) How is the variability in oxidation states of transition metals different from that of the p-block elements?
- (ii) Out of Cu^+ and Cu^{2+} , which ion is unstable in aqueous solution and why?
- (iii) Orange colour of $Cr_2O_7^{2-}$ ion changes to yellow when treated with an alkali. Why?
- (b) Chemistry of actinoids is complicated as compared to lanthanoids. Give two reasons.

Q.25. (a) An element has atomic mass 93 g mol^{-1} and density 11.5 g cm^{-3} . If the edge length of its unit cell is 300 pm, identify the type of unit cell. 5 marks

- (b) Write any two differences between amorphous solids and crystalline solids.

OR


(a) Calculate the number of unit cells in 8.1 g of aluminium if it crystallizes in a f.c.c. structure. (Atomic mass of Al = 27 g mol^{-1})

(b) Give reasons:


- (i) In stoichiometric defects, NaCl exhibits Schottky defect and not Frenkel defect.
- (ii) Silicon on doping with Phosphorous forms n-type semiconductor.
- (iii) Ferrimagnetic substances show better magnetism than antiferromagnetic substances.

Q.26. (a) Write the product(s) in the following reactions: 5 marks

(i)



(ii)



(iii) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_2 - \text{OH} \xrightarrow{\text{PCC}} ?$

(b) Give simple chemical tests to distinguish between the following pairs of compounds:

(i) Ethanol and Phenol

(ii) Propanol and 2-methylpropan-2-ol

OR

(a) Write the formula of reagents used in the following reactions:

(i) Bromination of phenol to 2,4,6-tribromophenol

(ii) Hydroboration of propene and then oxidation to propanol.

(b) Arrange the following compound groups in the increasing order of their property indicated:

(i) p-nitrophenol, ethanol, phenol (acidic character)

(ii) Propanol, Propane, Propanal (boiling point)

(c) Write the mechanism (using curved arrow notation) of the following reaction :



Buy solution: <http://www.4ono.com/cbse-12th-chemistry-solved-previous-year-papers/>