

# ICSE Paper 2005

## **PHYSICS**

## Section-I (40 marks)

Compulsory: (Attempt all questions from this Section)

	Compulsory: (Attempt all questions from this Bection)	
hie	estion 1.	. 44.
a)	Which physical quantity does the electron bott measures flow is it retailed to	[2]
	a T unit of this auantity?	
b)	What should the angle between force and displacement be to get the:	ro1
130	6) minimum work. (11) Maximum work:	[2]
c)	at to Newton's second law of motion.	[2]
d)	The work done by the heart is I jouie per beat. Calculate the power of the h	eart,
	:: Leate 79 times in one minute.	[4]
e)	Mention two properties of a wave: one property which varies and the	other
035	which remains constant when the wave passes from one medium to another	•
		[2]
ne	swer.	
a)	Energy is measure in electron volt.	
	$1 \text{ eV} = 1.6 \times 10^{-19} \text{ Joule.}$	
ы	The angle between force and displacement for:	
W)	(i) minimum work—Angle should be 90°.	
	(ii) maximum work—Angle should be 0°.	
(d)	SE 20 CONTROL OF THE THE PART OF THE PART	
u		
	Therefore in 72 beats = 72 joule Workdone	
	$Power = \frac{VOINGER}{Time}$	
	$=\frac{72 \text{ joule}}{60 \text{ sec.}}=\frac{6}{5}$	
	= 1·2 Watt	Ans.
(م)		
(E)	When a wave passes from one medium to another:	
	(i) Wavelength or velocity varies (changes).	
	(ii) Frequency remains constant.	
	estion 2.	F01
(a)	Explain briefly what causes the twinkling of stars at night.	[2]
(р)	State two advantages of an aneroid barometer.**	[2]
(e)	Explain why a gas bubble released at the bottom of a lake grows in size	as it
2000	rises to the surface of the lake.**	[2]
(d)	) What is meant by the statement, the critical angle for diamond is 24 $^\circ$ : $\epsilon$	low is
	the critical angle of a material related to its refractive index?	[2]
(e)	A block of wood of volume 25 cm3 floats in water with 20 cm3 of its v	оште
	immersed. Calculate:**	
2000	<ol> <li>The density and</li> <li>The weight of the block of wood.</li> </ol>	[2]

Answer has not given due to out of present syllabus.

## 704 | ICSE Last 10 Years Solved Papers

#### Answer.

(a) The phenomenon of the change in direction of a ray of light, in going from on medium to another is called refraction.

Twinkling of stars due to refraction of light by the atmosphere.

(d) It means that a ray of light is incident on a diamond air interface at an angumere than 24°, it would be totally internally reflected back.

The relation between critical angle and refractive index is

$$\mu = \frac{1}{\sin i_c}$$

### Question 3.

- (a) Name any two electromagnetic waves which have a frequency higher than that of violet light. State one use of each.
- (b) State two ways by which the frequency of transverse vibrations of a stretched string can be decreased.
- (c) Why does the temperature of the surroundings start falling when the ice of frozen lake start melting?
- (d) Four resistances of 2.0Ω each arc joined end to end, to form a square ABCD Calculate the equivalent resistance of the combination between any two adjacent corners.
- (e) In a three-pin plug, why is the earth pin made longer and thicker than the other two pins?


#### Answer.

- (a) The frequency of X-rays and gama-rays is higher than violet light. X-rays are used in X-ray crystallography. Gamma rays are used in radiotherapy.
- (b) The frequency of transverse vibration can be decreased by:
  - (i) Increasing the length (T and m keeping constant)
  - (ii) Decreasing the tension (l and m keeping constant)
- (c) When ice converts into water it needs 80 cal. heat per gram. Therefore, when frozen lake starts melting it withdraw a large amount of heat from the surrounding. Therefore temperature of surrounding starts falling.
- (d) Equivalent resistance between A and B:

$$\frac{1}{R} = \frac{1}{R_{AB}} + \frac{1}{R_{BC} + R_{CD}} + \frac{1}{R_{AD}}$$

$$\frac{1}{R} = \frac{1}{2} + \frac{1}{2 + 2 + 2} = \frac{1}{2} + \frac{1}{6}$$

$$\frac{1}{R} = \frac{3 + 1}{6} = \frac{4}{6}$$



Ans

$$R = \frac{6}{4} = 1.5 \Omega \text{ (ohm)}.$$

- (e) The earth pin is made longer and thicker because:
  - (i) To make save that the body of appliance can a good contact with the earth.
  - (ii) To avoid the chance of confusion between earth pin and other two pins.



## Question 4.

- (a) State the energy change which takes place when a magnet is moved inside a coil having a galvanometer at its ends. Name this phenomenon. [2]
- (b) Draw a labelled diagram of an A.C. generator.


- (c) Calculate the heat energy that will be released when 5.0 kg of steam at 100°C condenses to form water at 100°C. Express your answer in S.I. unit. (Specific latent heat of vaporization of steam is 2268 kJ/kg.)
- (d) How many alpha and beta particles are emitted when Uranium nucleus  $^{238}_{92}U$ decays to Lead 82 Pb ?
- (e) With the help of an equation, state the mechanism of energy production in a nuclear fusion reaction. \*\* [2]

## Answer.

(a) When a magnet is moved inside a coil (galvanometer at its ends), mechanical energy converts into electrical energy.

The phenomenon is known as electromagnetic induction.

**(b)** 



(c) Given: m = 5 kg, specific latent heat of steam = 2268 kJ/kg.

We know that,

$$Q = mL$$
  
=  $5 \times 2268 = 11340 \text{ kJ}$   
=  $11.34 \times 10^6 \text{ joule.}$   
 $38 \longrightarrow 82 \text{Pb}^{206}$ 

Ans.

(d)

Change in Atomic no.  $\approx 92 - 82 = 10$ 

To decrease in mass by 32 units 8  $\alpha$ -particle should be emitted

To decrease in Atomic no. 8  $\alpha$ -particle reduced the Atomic no.  $8 \times 2 = 16$  unit. But atomic no. is reduced by 10.

∴ 6-Beta particle should be emitted.

$$-16+6=-10$$

No. of  $\alpha$ -particle = 8

No. of 
$$\beta$$
-particle = 6

Ans.

Answer has not given due to out of present syllabus.



## 708 | ICSE Last 10 Years Solved Papers

(iii) If a ray of light passes from medium I to medium II without any change of direction, what can be said about the refractive indices of these media (angle i is not 0)?

### Answer.

- (a) (i) The magnification is 1.
  - (ii) The image real, inverted and of the same sized as the object.
  - (iii) The distance between the object and the optical centre of the lens equal the focal length of the lense.
- (b) (i) No, the letter of each colour has different (slightly) height.
  - (2) The letter V (violet) is raised more.

Because 'µ' is maximum for violet colour, therefore, it raised more.

- (ii) The colour of an object is green.
- (c) (i) When a ray of light passes from one medium to the other medium, it changes its path. This phenomenon is called Refraction of light. In passing from a rarer to a denser medium, it bends towards the normal passing from a denser to a rarer medium, it bends away from the normal
  - (ii) (1) Refractive Index  $(_1\mu_2) = \frac{Velocity\ of\ light\ in\ first\ medium}{Velocity\ of\ light\ in\ second\ medium}$ 
    - (2) Refractive index  $(_1\mu_2) = \frac{\sin i}{\sin r}$
  - (iii) If light ray is passing through another medium without changing its part even when, i is not 0, the refractive index of two medium must be equal.

## Question 8.


- (a) A radar is able to detect the reflected waves from an enemy aeroplane, after time interval of 0.02 milliseconds. If the velocity of the waves is 3 × 10<sup>8</sup> ms, calculate the distance of the plane from the radar.
- (b) A piece of ice is heated at a constant rate. The variation of temperature 110°C with 100°C heat input is shown in the graph

below:
(i) What are represented by AB and

CD?

(ii) What conclusion can you draw regarding the nature of ice from - 10°C

the above graph? [3]



- (c) If there is no heat loss to the surroundings, the heat released by the condessation of m<sub>1</sub> g of steam at 100°C into water at 100°C can be used to convert more of ice at 0°C into water at 0°C.
  - (i) *Find*:
    - (1) The heat lost by steam in terms of m1.
    - (2) The heat gained by ice in terms of m2.
  - (ii) Form a heat equation to find the ratio of m<sub>2</sub>: m<sub>1</sub>. (Specific latent heat of vaporization of steam = 2268 kJ/kg; Specific latent heat of fusion of ice = 336 kJ/kg. Specific heat capacity of water = 4200 J/kg°C.



Answer.

(a) Let x be the distance of plane from the Radar then

$$x + x = 2x = \text{velocity} \times \text{time}$$

$$2x = 3 \times 10^8 \times \cdot 02 \times 10^{-3}$$

$$= 6 \times 10^3 \text{ m} = 6 \text{ km}.$$

Ans.

- → represents the time interval in which ice changes into water at
  - ---- represents the time interval in which water converts into · vapour at 100°C.
  - (ii) The initial temperature of ice is ~10°C, it can possible only by addition of some salt or under high pressure.
- (1) The heat lost by the steam =  $m_1 L_a$  $(L_s = Specific latent heat of steam)$ 
  - (2) Heat gained by the ice =  $m_2 L_i$ (L; = Specific latent heat of ice)
  - Heat given by steam = Heat taken by ice (ii)

$$m_1 L_s = m_2 L_i$$

$$\frac{m_2}{m_1} = \frac{L_s}{L_i} = \frac{2268}{336} = \frac{27}{4}$$

$$\mathbf{m_2} : \mathbf{m_1} = 27 : 4$$

Ans.

Question 9.

(a) An electrical appliance is rated 1500 W, 250 V. This appliance is connected to 250 V mains.

Calculate:

- (i) the current drawn,
- (ii) the electrical energy consumed in 60 hours,


(iii) the cost of electrical energy consumed at Rs. 2.50 per kWh. (b) (i) State the function of a split ring in a D.C. motor.

រោ

- (ii) Mention two reasons why a soft iron core is used within the coil of a moving coil galvanometer. [3]
- (c) In the figure given alongside, A, B and C are three ammeters. The ammeter B reads 0.5A. (All the ammeters have negligible resistance)

Calculate:

The readings in the ammeters A and



(ii) The total resistance of the circuit. Answer.

(a) (i) The current drawn

$$i = \frac{P}{V} = \frac{1500}{250} = 6A$$

Ans.

Electrical energy consume in 60 hours

$$W = P \times t = 1500 \times 60 = 90 \text{ kWh}$$
  
= 1500 \times 60 \times 60 \times 60  
= 3240 \times 10<sup>5</sup> joule.

Ans.



## 710 | ICSE Last 10 Years Solved Papers

(iii)  $Cost = 90 \times 2.50$ = Rs. 225.

- (b) (i) The function of split rings in d.c. motor to flow the current in armature coil in such a way that it always keep on rotating in the same manner.
  - (ii) Soft iron core is used in moving coil galvanometer, because
    - (a) To make the field 'radial'.
    - (b) To concentrate the magnetic lines of force.

(c) (i) 
$$\frac{\text{Ammeter 'C' Reading}}{\text{Ammeter 'B' Reading}} = \frac{6}{3} = 2$$

$$\text{Reading of 'C'} = 2 \times .5$$

$$= 1.0 \text{ A}$$

$$\text{Reading of 'A'} = (1.0 + 0.5)$$

$$= 1.5 \text{ A}$$

(ii) The total resistance of the circuit

R = 2 + Resistance conneted in parallel

Ang

[3]

$$\frac{1}{R} = \left(\frac{1}{3} + \frac{1}{6}\right) = \left(\frac{2+1}{6}\right) = \frac{3}{6} \implies R = 2\Omega$$

$$R = 2+2$$

$$= 4\Omega.$$

Question 10.

- (a) State the functions of the following in a nuclear reactor: \*\*
  - (i) Moderator,
  - (ii) Control rods,
  - (iii) Coolant.
- (b) (i) Mention two important precautions that should be taken while handling radioactive materials.
  - (ii) State one use of radio-isotopes.
- (c) (i) Draw a labelled diagram of a hot cathode ray tube.
  - (ii) Why are materials of low work function preferred as thermionic cathods materials?
  - (iii) Write an equation to show the fission of a nucleus of  $U^{235}$  with the production of three neutrons. \*\*

### Answer.


- (b) (i) While using radioactive material it is essential that
  - (1) put them in a thick lead box.
  - (2) wear special protective (led coated) clothes and gloves on the hand.
  - (ii) Radioisotopes are used in medical diagnosis and in radiotheraphy.

<sup>\*\*</sup> Answer has not given due to out of present syllabus.

Physics, 2005 | 711

(c) (i) Please see given below figure.

1.35



(ii) To get more electron at relatively low temperature, we use the material of low work function.



